# East Carolina Gastroenterology, PA Jacksonville, NC 28546 Peter A. Eweje, M. D. 910 353 6158

#### **INSTRUCTIONS FOR MAGNESIUM CITRATE PREP**

- Your procedure is scheduled at **our Endoscopy Center/OMH** on \_\_\_\_\_\_at \_\_\_\_\_at \_\_\_\_at \_\_\_\_\_at \_\_\_\_at \_\_\_\_\_at \_\_\_\_\_\_at \_\_\_\_\_\_at \_\_\_\_\_\_at \_\_\_\_\_\_at \_\_\_\_\_\_\_at \_\_\_\_\_\_at \_\_\_\_\_\_\_at
- Obtain your bowel prep products from your pharmacy and buy Gas-X from over-the counter
- Please stop taking your aspirin, celebrex, Plavix, motrin, aleve, and other ibuprofen containing medications five days before your test, unless we instruct you otherwise.
- If your test is <u>scheduled for a morning DO NOT take any of your regular a.m</u> <u>medications(except inhalers and medicines for seizures)</u> on the day of your test. You <u>MAY</u> <u>TAKE your regular a.m medications if your test is scheduled for the afternoon.</u> Continue to take your inhalers and seizure medications as prescribed by your physician. Bring your medication bottles along with you. Do not hesitate to call us with questions about your medications.
- You will need specific instructions if you take Coumadin/Warfarin, Pradaxa, Xarelto, Eliquis, Savaysa or other anticoagulants. Please let us know if you do take any of these medications. You may need to be switched to a blood thinning shot.
- **Do not take your diabetic pills** the evening before and the morning of your test. You should only **take half the usual dose** of your **Insulin shot** the day before your test, and none on the morning of your test.
- In preparation for your procedure try to **eat less of foods high in fiber** for at least **5 days** before the procedure. Some foods to avoid include: bran, whole grain cereals, all beans, raw fruits and vegetables, potato skins, nuts, raisins, corn, seeds and popcorn.

### THE DAY BEFORE THE PROCEDURE( )

- Have a regular breakfast before 10am on \_\_\_\_\_. Then start CLEAR LIQUIDS, and continue until two hours before your test. See attached list for clear liquid diet suggestions. NO SOLID FOOD AFTER 10AM THE DAY BEFORE YOUR TEST
- At 4pm take 4 tablets of Dulcolax with two glasses of water.
- At 5pm drink the full bottle of Magnesium Citrate with 4 full glasses of water
- Take 2 Gas-X tablet with 8oz of clear liquid at 8pm.

### THE DAY OF THE PROCEDURE( )

- 5 hours(\_\_\_\_\_) before your test take 4 tablets of Dulcolax with two glasses of water.
- 4 hours(\_\_\_\_\_) before your test drink the full bottle of Magnesium Citrate with 4 full glasses of water
- Take 2 Gas-X tablet after drinking the Magnesium Citrate.
- You may continue with a clear liquid diet until 2 hours before your test.
- EAT <u>NO</u> SOLID FOOD.
- DO NOT DRINK OR EAT ANYTHING FOR 2 HOURS BEFORE YOUR TEST.
- You may apply a petroleum based ointment or diaper rash ointment to the rectal area if you experience discomfort.
- YOU WILL NEED A <u>**RESPONSIBLE ADULT**</u> TO DRIVE YOU HOME AFTER YOUR PROCEDURE. PLAN ON TAKING THE WHOLE DAY OFF OF WORK/SCHOOL THE DAY OF YOUR PROCEDURE.
- Please call the office at **910 353-6158** if you have any questions regarding your prep or procedure.

## <u>PLEASE FOLLOW THE INSTRUCTIONS ABOVE TO ENSURE YOUR COLON IS</u> <u>CLEAN. WE MAY NEED TO REPEAT THE TEST IF YOUR COLON IS NOT CLEAN</u>

PLEASE VISIT THE "PATIENT EDUCATION" SECTION ON OUR WEBSITE www.eastcarolinagastro.com TO LEARN MORE ABOUT YOUR PROCEDURE